



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 623 607 A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 94106320.8

(22) Date of filing: 22.04.94

(51) Int. Cl.⁵: **C07D 307/80, C07D 307/81, C07D 407/08, C07D 405/10, C07D 405/12, C07D 407/10, C07D 407/12, C07D 413/12, C07D 413/14, A61K 31/34, A61K 31/35, A61K 31/41, A61K 31/44, A61K 31/445, A61K 31/47, A61K 31/535, C07D 405/08**

(30) Priority: 06.05.93 GB 9309324

(43) Date of publication of application:
09.11.94 Bulletin 94/45

(94) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL PT SE

(71) Applicant: **BAYER AG**
D-51368 Leverkusen (DE)

(72) Inventor: Fischer, Rüdiger, Dr.
Pahlkestrasse 5
D-42115 Wuppertal (DE)
Inventor: Bräunlich, Gabriele, Dr.
Claudiusweg 9
D-42115 Wuppertal (DE)
Inventor: Mohrs, Klaus-Helmut, Dr.
Wildsteig 24
D-42113 Wuppertal (DE)

Inventor: **Hanko, Rudolf, Dr.**
Waldsaum 25

D-45134 Essen (DE)

Inventor: **Butler-Ransohoff, John-Edward, Dr.**
Pahlkestrasse 5

D-42115 Wuppertal (DE)

Inventor: **Es-Sayed, Mazen, Dr.**
Falkenberg 39

D-42113 Wuppertal (DE)

Inventor: **Sturton, Graham, Dr.**

184 Windsor Road
Bray Maidenhead SL6 2DW (GB)

Inventor: **Tudhope, Steve, Dr.**

47 Kentons Lane
Windsor Berkshire SL4 4JH (GB)

Inventor: **Abram, Trevor, Dr.**


214 Marlow Bottom
Marlow Buckinghamshire (GB)

Inventor: **McDonald-Gibson, Wendy J., Dr.**

2 Cottesmore Lane
Ewelme Wallingford, Oxford (GB)

(54) **Benzofuranyl-and-thiophenyl-alkanecarboxylic acid derivatives.**

(57) The benzofuranyl- and -thiophenyl-alkane-carboxylic acid derivatives of the formula (I)



wherein R₁, R₂, R₃, T, V, and W are as defined in Claim 1 are prepared by cyclisation of hydroxy acetophenones and related compounds or by Wittig-reaction of benzofuranyl aldehydes. The compounds can be

used to prepare medicaments showing antiinflammatory activity.

[illegible][illegible]

the 1990s, the number of people in the world who are illiterate has increased from 1.2 billion to 1.5 billion. The number of illiterate people in the world is projected to reach 1.7 billion by the year 2015. The number of illiterate people in the world is projected to reach 1.7 billion by the year 2015. The number of illiterate people in the world is projected to reach 1.7 billion by the year 2015.


The invention relates to benzofuranyl- and -thiophenyl-alkanecarboxylic acids derivatives, processes for their preparation and their use in medicaments.

It is known that the NADPH oxidase of phagocytes is the physiological source to the superoxide radical anion and reactive oxygen species derived therefrom which are important in the defence against pathogens. Uncontrolled formation leads to tissue damage in inflammatory processes. It is additionally known that elevation of phagocyte cyclic AMP leads to inhibition of oxygen radical production and that this cell function is more sensitive than others such as aggregation or enzyme release (cf. *Imb. Arch. Allergy Immunol.*, vol. 97: pp 194-199, 1992).

Benzofuran- and benzothiophene derivatives having lipoxygenase-inhibiting action are described in the publication EP 146 243.

Surprisingly it was found that compounds given by the general formula (I) inhibited oxygen radical formation and elevated cellular cyclic AMP levels probably by inhibition of phagocyte phosphodiesterase activity.

The invention relates to benzofuranyl- and -thiophenyl-alkanecarboxylic acids derivatives of the general formula (I)



in which

R^1 and R^2 are identical or different and represent hydrogen, halogen, carboxyl, cyano, nitro, trifluoromethyl or a group of a formula -OR⁴, -SR⁵ or -NR⁶R⁷,


in which

R^5 denotes hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms, R^4 , R^6 and R^7 are identical or different and denote hydrogen, cycloalkyl having 3 to 6 carbon atoms, or a 5 to 7-membered saturated or unsaturated heterocycle having up to 3 heteroatoms from the series comprising N, S and O, which are optionally substituted by identical or different substituents from the series comprising halogen, cyano, nitro or by straight-chain or branched alkyl or alkoxyacetyl each having up to 6 carbon atoms, or denote a residue of formula




or

denote straight-chain or branched alkyl or alkenyl each having up to 8 carbon atoms, and each of which is optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising trifluoromethyl, difluoromethyl, halogen, cyano, carboxy, hydroxy, straight-chain or branched alkoxy, alkoxyacetyl or acyl each having up to 6 carbon atoms or by a 5- to 7-membered saturated or unsaturated heterocycle having up to 3 heteroatoms from the series comprising N, S and O and to which an aromatic ring can be fused, or by N-methyl-substituted imidazolyl or by a residue of formula



- or by phenyl, where in all rings are optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising nitro, halogen, carboxy or straight-chain or branched alkyl or alkoxycarbonyl each having up to 6 carbon atoms,
- 5 or alkyl or alkenylene are substituted by a group of formula $-\text{CO}-\text{NR}^8\text{R}^9$ in which
- R^8 and R^9 are identical or different and denote phenyl, adamantyl, cycloalkyl having up to 3 to 7 carbon atoms, benzyl, formyl, hydrogen, straight-chain or branched alkyl or alkenyl each having up to 6 carbon atoms and which are optionally substituted by carboxy,
- 10 hydroxy or straight-chain or branched alkoxycarbonyl up to 6 carbon atoms or
- R^8 and R^9 together with the nitrogen atom form a 5 to 7 membered saturated or unsaturated heterocycle,
- or
- 15 R^4 denotes a protecting group of a hydroxyl group, difluoromethyl or a group of a formula $-\text{SO}_2-\text{X}$ in which
- X denotes trifluoromethyl, phenyl or straight-chain or branched alkyl having up to 6 carbon atoms,
- 20 T represents an oxygen or sulfur atom
- V represents a straight-chain or branched alkylene or alkenylene chain each having 2 to 8 carbon atoms,
- W represents cyano, tetrazolyl or a group of a formula $-\text{CO}-\text{R}^{10}$, $-\text{CO}-\text{NR}^{11}\text{R}^{12}$, $-\text{CONR}^{13}-\text{SO}_2-\text{R}^{14}$ or $\text{PO}(\text{OR}^{15})(\text{OR}^{16})$, or a residue of the formula



- 25 in which
- R^{10} denotes hydroxyl, cycloalkyloxy having up to 3 to 7 carbon atoms or straight-chain or branched alkoxy having up to 8 carbon atoms,
- 35 R^{11} , R^{12} and R^{13} are identical or different and denote hydrogen, phenyl or straight-chain or branched alkyl or acyl each having up to 6 carbon atoms and which are optionally substituted by hydroxyl,
- or
- R^{11} denotes hydrogen and
- 40 R^{12} denotes hydroxyl
- or
- R^{11} and R^{12} together with the nitrogen atom form a 5- or 6-membered saturated heterocycle,
- R^{14} denotes straight-chain or branched alkyl having up to 6 carbon atoms, which is optionally substituted by phenyl or trifluoromethyl, or denotes phenyl, which is optionally substituted by substituents from the series comprising halogen, cyano,
- 45 nitro or by straight-chain or branched alkyl having up to 6 carbon atoms,
- R^{15} and R^{16} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms,
- R^3 represents phenyl, which is monosubstituted to trisubstituted by identical or different substituents from the series comprising hydroxyl, halogen, nitro, tetrazolyl,
- 50 trifluoromethoxy, difluoromethoxy, trifluoromethyl, difluoromethyl, cyano, carboxy, straight-chain or branched alkyl, alkylthio, alkoxy, alkoxycarbonyl or acyl each having up to 8 carbon atoms or by a group of formula $-\text{NR}^{17}\text{R}^{18}$, $-(\text{O})_a-\text{SO}_2-\text{R}^{19}$ or $-\text{SO}_2-\text{NR}^{20}\text{R}^{21}$
- 55 in which
- a denotes a number 0 or 1,
- R^{17} and R^{18} have the meaning shown above for R^{11} and R^{12} and are identical to the latter or

- different from the latter,
or
R¹⁷ denotes hydrogen
and
5 R¹⁸ denotes straight-chain or branched acyl having up to 6 carbon atoms
and
R¹⁹ has the above mentioned meaning of R¹⁴ and is identical to the latter or different
from the latter.
R²⁰ and R²¹ have the above mentioned meaning of R¹¹ and R¹² and are identical to the latter or
10 different from the latter

and salts thereof.

The benzofuranyl- and -thiophenyl-alkanecarboxylic acids derivatives according to the invention can also be present in the form of their salts. In general, salts with organic or inorganic bases or acids may be mentioned here.

- 15 Physiologically acceptable salts are preferred in the context of the present invention. Physiologically acceptable salts of the benzofuranyl- and -thiophenyl-alkanecarboxylic acids derivatives can be metal or ammonium salts of the substances according to the invention, which contain a free carboxylic group. Those which are particularly preferred are, for example, sodium, potassium, magnesium or calcium salts, and also ammonium salts which are derived from ammonia, or organic amines, such as, for example, ethylamine, di-
20 or triethylamine, di- or triethanolamine, dicyclohexylamine, dimethylaminoethanol, arginine, lysine or ethylenediamine.

- Physiologically acceptable salts can also be salts of the compounds according to the invention with inorganic or organic acids. Preferred salts here are those with inorganic acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid or sulphuric acid, or salts with organic carboxylic or
25 sulphonic acids such as, for example, acetic acid, maleic acid, fumaric acid, malic acid, citric acid, tartaric acid, ethanesulphonic acid, benzenesulphonic acid, toluenesulphonic acid or naphthalenedisulphonic acid.

- The compounds according to the invention can exist in stereoisomeric forms which either behave as image and mirror image (enantiomers), or which do not behave as image and mirror image (diastereomers). The invention relates both to the antipodes and to the racemate forms, as well as the diastereomer
30 mixtures. The racemate forms, like the diastereomers, can be separated into the stereoisomerically uniform constituents in a known manner.

Hydroxyl protective group in the context of the above-mentioned definition in general represents a protective group from the series comprising: trimethylsilyl, tert.butyl-dimethylsilyl, benzyl, 4-nitrobenzyl, 4-methoxybenzyl, acetyl, tetrahydropyranyl and benzoyl.

- 35 Heterocycle in general represents a 5- to 7-membered, preferably 5- to 6-membered, saturated or unsaturated ring which can contain up to three oxygen, sulphur and/or nitrogen atoms as heteroatoms and to which further aromatic ring can be fused.

5- and 6-membered rings having an oxygen, sulphur and/or up to two nitrogen atoms are preferred, which may also be fused to benzene.

- 40 The following are mentioned as preferred: thienyl, furyl, pyrrolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, quinolyl, isoquinolyl, quinazolyl, quinoxazolyl, cinnolyl, thiazolyl, benzothiazolyl, isothiazolyl, benzisothiazolyl, oxazolyl, benzoxazolyl, isoxazolyl, imidazolyl, benzimidazolyl, indolyl, morpholyl, pyrrolidinyl, piperidyl or piperazinyl.

Preferred compounds of the general formula (I) are those

- 45 in which
R¹ and R² are identical or different and represent hydrogen, fluorine, chlorine, bromine, nitro, trifluoromethyl or a group of a formula -OR⁴, -SR⁵ or -NR⁶R⁷,
in which
R⁶ denotes hydrogen or straight-chain or branched alkyl having up to 3 carbon atoms,
50 R⁴, R⁵ and R⁷ are identical or different and denote cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, chinolyl, pyridyl, imidazolyl, 1,3-thiazolyl or thienyl, which are optionally substituted by identical or different substituents from the series comprising fluorine, chlorine, bromine, iodine, cyano, nitro or by straight-chain or branched alkyl or alkoxy-carbonyl
each having up to 5 carbon atoms or
55 denote a residue of formula



5


or

denote straight-chain or branched alkyl or alkenylen each having up to 6 carbon atoms, and each of which is optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising trifluoromethyl, difluoromethyl, fluorine, chlorine, bromine, iodine, cyano, carboxy, hydroxy, straight-chain or branched alkoxy, alkoxycarbonyl or acyl each having up to 5 carbon atoms or by chinolyl, pyridyl, pyrazolyl, 1,3-thiadiazolyl; thienyl, imidazolyl or N-methyl-substituted imidazolyl, and to which an aromatic ring can be fused,


10

or by a residue of formula

15



20



25

or by phenyl; where in all rings are optionally monosubstituted to disubstituted by identical or different substituents from the series comprising nitro, fluorine, chlorine, bromine, iodine, carboxy or straight-chain or branched alkyl or alkoxycarbonyl each having up to 5 carbon atoms;

30

or alkyl or alkenylen are substituted by a group of formula $-\text{CO}-\text{NR}^8\text{R}^9$

in which

R^8 and R^9

are identical or different and denote phenyl, adamantyl, cyclopropyl, cyclopentyl, benzyl, formyl, hydrogen, straight-chain or branched alkyl or alkenyl each having up to 5 carbon atoms, which are optionally substituted by carboxy, hydroxy or straight-chain or branched alkoxycarbonyl up to 4 carbon atoms,

35

or

R^8 and R^9

together with the nitrogen atom form a morpholinyl, piperidinyl, piperazinyl or a pyrrolidinyl ring,

40

or

R^4

denotes acetyl, benzyl, tetrahydrofluorenyl, difluoromethyl or a group of a formula $-\text{SO}_2-\text{X}$, in which

45

X

denotes trifluoromethyl, phenyl or methyl,

T

represents an oxygen or sulfur atom


V

represents a straight-chain or branched alkylene or alkenylene chain each having 2 to 6 carbon atoms,

W

represents cyano, tetrazolyl or a group of a formula $-\text{CO}-\text{R}^{10}$, $-\text{CO}-\text{NR}^{11}\text{R}^{12}$, $-\text{CONR}^{13}-\text{SO}_2-\text{R}^{14}$ or $\text{PO}(\text{OR}^{15})(\text{OR}^{16})$, or a residue of the formula

50



55

in which

- R¹⁰ denotes hydroxyl, cyclopropyloxy, cyclopentyloxy, cyclohexyloxy or straight-chain or branched alkoxy having up to 6 carbon atoms,
 R¹¹, R¹² and R¹³ are identical or different and denote hydrogen, phenyl or straight-chain or branched alkyl or acyl each having up to 4 carbon atoms and which are optionally substituted by hydroxyl,
 or
 R¹¹ denotes hydrogen
 and
 R¹² denotes hydroxyl
 or
 R¹¹ and R¹² together with the nitrogen atom form a pyrrolidiny, piperidiny ring or a morpholiny,
 R¹⁴ denotes straight-chain or branched alkyl having up to 5 carbon atoms, which is optionally substituted by phenyl or trifluoromethyl, or denotes phenyl, which is optionally substituted by substituents from the series comprising fluorine, chlorine, bromine, iodine, cyano, nitro or by straight-chain or branched alkyl having up to 4 carbon atoms,
 R¹⁵ and R¹⁶ are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms,
 R³ represents phenyl, which is monosubstituted to trisubstituted by identical or different substituents from the series comprising hydroxyl, fluorine, chlorine, bromine, iodine, nitro, trifluoromethoxy, difluoromethoxy, trifluoromethyl, difluoromethyl, cyano, carboxy, tetrazolyl, straight-chain or branched alkyl, alkylthio, alkoxy, alkoxycarbonyl or acyl each having up to 6 carbon atoms or by a group of formula -NR¹⁷R¹⁸, -(O)_a-SO₂-R¹⁹ or -SO₂NR²⁰R²¹,
 in which
 a denotes a number 0 or 1,
 R¹⁷ and R¹⁸ have the meaning shown above for R¹¹ and R¹² and are identical to the latter or different from the latter,
 or
 R¹⁷ denotes hydrogen
 and
 R¹⁸ denotes straight-chain or branched acyl having up to 6 carbon atoms
 and
 R¹⁹ has the above mentioned meaning of R¹⁴ and is identical to the latter or different from the latter.
 and
 R²⁰ and R²¹ have the above mentioned meaning of R¹¹ and R¹² and are identical to the latter or different from the latter,

and salts thereof.



Particularly preferred compounds of the general formula (I) are those

in which

- R¹ denotes hydrogen,
 R² represents fluorine, chlorine, bromine, nitro, trifluoromethyl or a group of a formula -OR⁴ or -NR⁵R⁷,
 in which
 R⁴ denotes a group of a formula -SO₂X,
 in which
 X denotes trifluoromethyl, phenyl, methyl or difluoromethyl,
 or
 R⁴ denotes hydrogen, tetrahydropyranyl, difluoromethyl, acetyl, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, chinolyl, pyridyl, imidazolyl or thienyl, which are optionally substituted by identical or different substituents from the series comprising fluorine, chlorine, bromine, cyano, nitro or by straight-chain or branched alkyl or alkoxycarbonyl each having up to 4 carbon atoms or
 denotes a residue of the formula



or
denotes straight-chain or branched alkyl or alkenylen each having up to 5 carbon atoms, and each of which is optionally monosubstituted to disubstituted by identical or different substituents from the series comprising trifluoromethyl, difluoromethyl, fluorine, chlorine, bromine, cyano, carboxy, hydroxy, straight-chain or branched alkoxy, alkoxycarbonyl or acyl each having up to 5 carbon atoms or by chinolyl, pyridyl, imidazolyl or N-methyl substituted imidazolyl, and to which an aromatic ring can be fused, or by a residue of formula



or by phenyl; wherein all rings are optionally monosubstituted to disubstituted by identical or different substituents from the series comprising nitro, fluorine, chlorine, bromine, carboxy or straight-chain or branched alkyl or alkoxycarbonyl each having up to 4 carbon atoms,

or alkyl or alkenylen are substituted by a group of formula $-CO-NR^8R^9$ in which

R^8 and R^9 are identical or different and denote phenyl, benzyl, adamantyl, cyclopropyl, cyclopentyl, formyl, hydrogen, straight-chain or branched alkyl or alkenyl each having up to 5 carbon atoms, which are optionally hydroxy, substituted by carboxy, hydroxy or straight-chain or branched alkoxycarbonyl up to 3 carbon atoms

or

R^8 and R^9 together with the nitrogen atom form a morpholinyl, piperidinyl, piperazinyl or a pyrrolidinyl ring,


R^6 denotes hydrogen, methyl or ethyl,

R^7 denotes hydrogen, methyl or ethyl,

T represents an oxygen or sulfur atom

V represents a straight-chain or branched alkylene or alkenylene chain each having 2 to 5 carbon atoms,

W represents cyano, tetrazolyl or a group of a formula $-CO-R^{10}$, $-CO-NR^{11}R^{12}$, $-CONR^{13}-SO_2-R^{14}$ or $PO(OR^{15})(OR^{15})$ or a residue of the formula



in which

R^{10} denotes hydroxyl, cyclopropyloxy, cyclopentyloxy, cyclohexyloxy or straight-chain or branched alkoxy having up to 5 carbon atoms,


R^{11} , R^{12} and R^{13} are identical or different and denote phenyl, hydrogen, straight-chain or branched alkyl or acyl each having up to 4 carbon atoms and which are optionally substituted

by hydroxyl,
or
R¹¹ denotes hydrogen
and
5 R¹² denotes hydroxyl
or
R¹¹ and R¹² together with the nitrogen atom form a pyrrolidiny, piperidiny or morpholiny ring,
R¹⁴ denotes straight-chain or branched alkyl having up to 4 carbon atoms, which is
10 optionally substituted by phenyl or trifluoromethyl, or denotes phenyl, which is
optionally substituted by substituents from the series comprising fluorine, chlorine,
bromine, cyano, nitro or by a straight-chain or branched alkyl having up to 3 carbon
atoms,
R¹⁵ and R¹⁶ are identical or different and represent hydrogen or straight-chain or branched alkyl
having up to 5 carbon atoms,
15 R³ represents phenyl, which is monosubstituted to trisubstituted by identical or different
substituents from the series comprising hydroxyl, fluorine, chlorine, bromine, nitro,
tetrazolyl, trifluoromethyl, difluoromethyl, difluoromethoxy, trifluoromethoxy, cyano,
carboxy, straight-chain or branched alkyl, alkylthio, alkoxy, alkoxycarbonyl or acyl
each having up to 5 carbon atoms or by a group of formula -NR¹⁷R¹⁸, -(O)_aSO₂-R¹⁹
20 or -SO₂-NR²⁰R²¹,
in which
a denotes a number 0 or 1,
R¹⁷ and R¹⁸ have the meaning shown above for R¹¹ and R¹² and are identical to the latter or
different from the latter,
25 or
R¹⁷ denotes hydrogen
and
R¹⁸ denotes straight-chain or branched acyl having up to 5 carbon atoms
and
30 R¹⁹ has the abovementioned meaning of R¹⁴ and is identical to the latter or different
from the latter,
R²⁰ and R²¹ have the above mentioned meaning of R¹¹ and R¹² and are identical to the latter or
different from the latter,

and salts thereof.

35 Processes for the preparation of the compounds of the general formula (I) have additionally been found,
characterised in that

[A] compounds of the general formula (II)



45 in which
R¹, T, W and V have the abovementioned meaning
R²² represents a group of formula -OR⁴


50 in which
R⁴ has the abovementioned meaning of R⁴, but does not represent hydrogen,
are reacted with compounds of the general formula (III)

R³-CO-CH₂-Y (III)

55 in which
R³ has the abovementioned meaning
and

Y represents a typical leaving group such as, for example, chlorine, bromine, iodine, tosylate or


mesylate, preferably bromine,
 in inert solvents and in the presence of a base under cyclisation by customary methods
 or
 [B] in the case, in which V represents alkenyl, compounds of the general formula (IV)



in which

R¹, R³, T and R²² have the abovementioned meaning,


first are converted by reaction with N-bromosuccinimide, in inert solvents and in the presence of a catalyst to the compounds of the general formula (V)



in which

R¹, R³, T and R²² have the abovementioned meaning,


and then by subsequent hydrolysis to compounds of the general formula (VI)



in which

R¹, R³, T and R²² have the abovementioned meaning,

which in a last step are reacted with compounds of the general formula (VII)



in which

R¹¹ and R¹² have the abovementioned meaning,


R²³ represents C₁-C₄-alkyl

in inert solvents and in the presence of a base,

and in the case of the free hydroxyl functions (R⁴ = H) the protective groups are removed by a customary method,

and in the case of the acids (R¹⁰ = OH), the esters are hydrolysed, and in the case of the variation of the esters (R¹⁰ ≠ OH) the acids are esterified with the appropriate alcohols in the presence of a catalyst according to a customary method,


and in the case of the amides and sulfonamides (R⁴/R⁵/R⁷ = -CONR⁸R⁹/W = CONR¹¹R¹² / -CONR¹³-SO₂R¹⁴), using amines of the formula (VIII) or sulfonamides of the formula (IX)



in which

[B]

5




10

N-bromosuccinimide

CCl₄, dibenzoylperoxide


15




20

NaHCO₃, DMSO

25




30



THF, BuLi

35



40

45

Suitable solvents are generally customary organic solvents which do not change under the reaction conditions. These preferably include ethers such as diethyl ether, dioxane, tetrahydrofuran or glycol dimethyl ether, acetone, dimethylsulfoxide, dimethylformamide, N-methylpyrrolidone, pyridyl, methylethyl- or methylisobutyl ketone.

Suitable bases are generally inorganic or organic bases. These preferably include alkali metal hydroxides such as, for example, sodium hydroxide, sodium hydrogen-carbonate or potassium hydroxide, alkaline earth metal hydroxides such as, for example, barium hydroxide, alkali metal carbonates such as sodium carbonate, potassium carbonate, alkaline earth metal carbonates such as calcium carbonate, or alkaline metal or alkaline earth metal alkoxides such as sodium methoxide or potassium methoxide, sodium ethoxide or potassium ethoxide or potassium tert.butoxide, or organic amines (trialkyl(C₁-C₆)amines) such as triethylamine, or heterocycles such as 1,4-diazabicyclo[2.2.2]octane (DABCO), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), or amides such as sodium amides, lithium butyl amid or butyllithium, pyridin or methylpiperidine. It is also possible to employ alkali metals, such as sodium or its hydrides such as sodium

hydride, as bases. Potassium carbonate, butyllithium and sodium hydrogencarbonate are preferred.

The base is employed in an amount from 1 mol to 10 mol, preferably from 1.0 mol to 2.1 mol, relative to 1 mol of the compounds of the general formula (III).

The reactions in general proceed in a temperature range from -70°C to $+100^{\circ}\text{C}$, preferably from -70°C to $+80^{\circ}\text{C}$ and at normal pressure.

The cyclisation in general proceeds in a temperature range from $+30^{\circ}\text{C}$ to $+180^{\circ}\text{C}$, preferably from $+60^{\circ}\text{C}$ to $+120^{\circ}\text{C}$ and at normal pressure.

The process according to the invention is in general carried out at normal pressure. However, it is also possible to carry out the process at elevated pressure or at reduced pressure (for example in a range from 0.5 to 5 bar).

Suitable solvents for the bromination are halogenohydrocarbons such as dichloromethane, trichloromethane, tetrachloromethane, dichloroethylene, trichloroethylene or chlorobenzene. Carbon tetrachloride is preferred for the bromination with N-bromosuccinimide, dichloromethane for the bromination with boron tribromide and glacial acetic acid for the bromination with hydrobromic acid.

Suitable catalysts for bromination are generally radical generators such as, for example, dibenzoyl peroxide or azobis-isobutyronitrile. Dibenzoyl peroxide is preferred.

The catalyst is employed in an amount from 0.001 mol to 0.2 mol, preferably from 0.1 mol to 0.05 mol, relative to 1 mol of the compounds of the general formula (IV).

The base is employed in an amount from 1 mol to 10 mol, preferably from 2.0 mol to 2.1 mol, relative to each to 1 mol of the compounds of the general formula (VII).

Bromination is generally carried out in a temperature range from -30°C to $+150^{\circ}\text{C}$, preferably from -20°C to $+50^{\circ}\text{C}$.

Bromination is generally carried out at normal pressure. However, it is also possible to carry out bromination at elevated pressure or at reduced pressure (for example in a range from 0.5 to 5 bar).

The process is in general carried out in a temperature range from $+10^{\circ}\text{C}$ to $+150^{\circ}\text{C}$, preferably from $+20^{\circ}\text{C}$ to $+100^{\circ}\text{C}$.

The process is generally carried out at normal pressure. However, it is also possible to carry out it at elevated pressure or at reduced pressure (for example in a range from 0.5 to 5 bar).

Suitable bases for the hydrolysis are the customary inorganic bases. These preferably include alkali metal hydroxides or alkaline earth metal hydroxides such as, for example, sodium hydroxide, potassium hydroxide or barium hydroxide, or alkali metal carbonates such as sodium carbonate or potassium carbonate or sodium hydrogen carbonate, or alkali metal alkoxides such as sodium methoxide, sodium ethoxide, potassium methoxide, potassium ethoxide or potassium tert.butoxide. Sodium hydroxide or potassium hydroxide are particularly preferably employed.

Suitable solvents for the hydrolysis are water or the organic solvents customary for hydrolysis. These preferably include alcohols such as methanol, ethanol, propanol, isopropanol or butanol, or ethers such as tetrahydrofuran or dioxane, or dimethylformamide, or dimethyl sulphoxide. Alcohols such as methanol, ethanol, propanol or isopropanol are particularly preferably used. It is also possible to employ mixtures of the solvents mentioned.

The hydrolysis can also be carried out with acids such as, for example, trifluoroacetic acid, acetic acid, hydrochloric acid, hydrobromic acid, methanesulphonic acid, sulphuric acid or perchloric acid, preferably with trifluoroacetic acid.

The hydrolysis is in general carried out in a temperature range from 0°C to $+180^{\circ}\text{C}$, preferably from $+20^{\circ}\text{C}$ to $+160^{\circ}\text{C}$.

In general, the hydrolysis is carried out at normal pressure. However, it is also possible to work at reduced pressure or at elevated pressure (for example from 0.5 to 5 bar).

When carrying out the hydrolysis, the base is in general employed in an amount from 1 to 3 mol, preferably from 1 to 1.5 mol, relative to 1 mol of the ester. Molar amounts of the reactants are particularly preferably used.

The amidation/sulphoamidation is in general carried out in one of the abovementioned solvents, preferably in dichloromethane. It may optionally proceed, starting from the free carboxylic acid, via an activated stage, for example via the corresponding acid halides, which can be prepared from the corresponding acids by reaction with thionyl chloride, phosphorus trichloride, phosphorus pentachloride, phosphorus tribromide or oxalyl chloride. Preferably, the activated stages are prepared from the corresponding acids using dicyclohexylcarbodiimide, N-(3-dimethylaminopropyl)-N'-thylcarbodiimide hydrochloride or carbonyldiimidazole and reacted in situ with the sulphonamides.

The amidation and the sulphoamidation are in general carried out in a temperature range from -20°C to $+80^{\circ}\text{C}$, preferably from -10°C to $+30^{\circ}\text{C}$ and at normal pressure.

In addition to the abovementioned bases, suitable bases for these reactions are preferably triethylamine and/or dimethylaminopyridine, DBU or DABCO.

The base is employed in an amount from 0.5 mol to 10 mol/l, preferably from 1 mol to 2 mol, relative to 1 mol of the appropriate ester or acid.


5 Acid-binding agents which can be employed for the sulfoamidation are alkali metal or alkaline earth metal carbonates such as sodium carbonate, potassium carbonate, alkali metal or alkaline earth metal hydroxides such as, for example, sodium hydroxide or potassium hydroxide, or organic bases such as pyridine, triethylamine, N-methyl-piperidine, or bicyclic amidines such as 1,5-diazabicyclo[3.4.0]-non-5-ene (DBN) or 1,5-diazabicyclo[3.4.0]undec-5-ene (DBU). Potassium carbonate is preferred.

10 Suitable dehydrating reagents are carbodiimides such as, for example, diisopropylcarbodiimide, dicyclohexylcarbodiimide or N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride or carbonyl compounds such as carbonyldiimidazole or 1,2-oxazolium compounds such as 2-ethyl-5-phenyl-1,2-oxazolium-3-sulphonate or propanephosphonic anhydride or isobutyl chloroformate or benzotriazolyloxy-tri-(dimethylamino)phosphonium hexafluorophosphate or diphenyl phosphoramidate or methanesulphonyl chloride, if appropriate in the presence of bases such as triethylamine or N-ethylmorpholine or N-methyl-piperidine or dicyclohexylcarbodiimide and N-hydroxysuccinimide.

The acid-binding agents and dehydrating reagents are in general employed in an amount from 0.5 to 3 mol, preferably from 1 mol to 1.5 mol, relative to 1 mol of the corresponding carboxylic acids.

The compounds of the general formula (II) and (III) are known or can be prepared by published methods.

20 The compound of the general formula (IV) are known in some cases and, for example, can be prepared by reacting compounds of the general formula (X)



in which

30 R₁, T and R₂₂ have the abovementioned meaning,

with compounds of the general formula (III) in one of the abovementioned solvents and bases, preferably acetone and potassium carbonate.

The base is employed in an amount from 1 mol to 10 mol, preferably from 1.0 mol to 2.1 mol, relative to 1 mol of the compounds of the general formula (X)

35 The reactions in general proceed in a temperature range from +30°C to +100°C, preferably from +40°C to +80°C and at normal pressure.

The compounds of the general formula (VIII), (IX) and (X) are known.

The compounds of the general formulas (V) and (VI) are known in some cases and can be prepared by the abovementioned processes.

40 The compounds of the general formula (VII) are known in some cases and can be prepared by customary methods.

The compounds according to the invention specifically inhibit the production of superoxide by polymorphonuclear leucocytes (PMN) without impairing other cell functions such as degranulation or aggregation. The inhibition was mediated by the elevation of cellular cAMP due to inhibition of the type IV

45 phosphodiesterase responsible for its degradation

They can therefore be employed in medicaments for controlling acute and chronic inflammatory processes.

The compounds according to the invention are preferably suitable for the treatment and prevention of acute and chronic inflammations of the airways, such as emphysema, alveolitis, shock lung, asthma, bronchitis, arteriosclerosis, arthrosis, inflammations of the gastro-intestinal tract and myocarditis. The compounds according to the invention are additionally suitable for reducing the damage to infarct tissue after reoxygenation. In this case the simultaneous administration of allopurinol to inhibit xanthine oxidase is of advantage. Combination therapy with superoxide dismutase is also of use.

55 Test description

1. Preparation of human PMN Blood was taken from healthy subjects by venous puncture and neutrophils were purified by dextran sedimentation and resuspended in the buffered medium.

2. Inhibition of FMLP-stimulated production of superoxide radical anions. Neutrophils ($2.5 \times 10^5 \text{ ml}^{-1}$) were mixed with cytochrome C (1.2 mg/ml) in the wells of a microtitre plate. Compounds according to the invention were added in dimethyl sulphoxid (DMSO). Compound concentration ranged from 2.5 nM to 10 μM , the DMSO concentration was 0.1% v/v in all wells. After addition of cytochalasin b ($5 \mu\text{g} \times \text{ml}^{-1}$) the plate was incubated for 5 min at 37°C . Neutrophils were then stimulated by addition of $4 \times 10^{-8} \text{ M}$ FMLP and superoxide generation measured as superoxide dismutase inhibitable reduction of cytochrome C by monitoring the OD_{550} in a Thermomax microtitre plate spectrophotometer. Initial rates were calculated using a Softmax kinetic calculation programme. Blank wells contained 200 units of superoxide dismutase.

The inhibition of superoxide production was calculated as follows:

$$\left[\frac{1 - ((R_x - R_b))}{(R_o - R_b)} \right] \cdot 100$$

R_x = Rate of the well containing the compound according to the invention.

R_o = Rate in the control well.

R_b = Rate in the superoxide dismutase containing blank well.

3. Measurement of PMN cyclic AMP concentration

The compounds according to the invention were incubated with 3.7×10^6 PMN for 5 min at 37°C before addition of $4 \times 10^{-8} \text{ M}$ FMLP. After 6 min protein was precipitated by the addition of 1% v/v conc. HCl in 96% v/v ethanol containing 0.1 mM EDTA. After centrifugation the ethanolic extracts were evaporated to dryness under N_2 and resuspended in 50 mM Tris/HCl pH 7.4 containing 4 mM EDTA. The cyclic AMP concentration in the extracts was determined using a cyclic AMP binding protein assay supplied by Amersham International plc. Cyclic AMP concentrations were expressed as percentage of vehicle containing control incubations.

Example No.	% elevation of [cAMP] at $1 \mu\text{M}$ (control 100)
30	394

4. Assay of PMN phosphodiesterase

PMN suspensions (10^7 cells/ml) were sonicated for 6 x 10 sec on ice. Aliquots (100 μl) were incubated for 5 min at 37°C with the compounds according to the invention or vehicle before the addition of ^3H -cAMP (1 mM and 200 nCi per incubation). After 20 min the reaction was stopped by heating at 100°C for 45 seconds. After cooling 100 mg of 5'-nucleotidase was added to each tube and the samples incubated for 15 min at 37°C . The conversion to ^3H -adenosine was determined by ion-exchange chromatography on Dowex AG-1x (chloride form) followed by liquid scintillation counting. Percentage inhibition was determined by comparison to vehicle containing controls.

5. Effect of intravenously administered compounds on the FMLP-induced skin oedema of guinea pigs

Guinea-pigs (600-800 g) were anaesthetized with pentobarbitone sodium (40 mg/kg, i.p.) and injected (i.v.) with a 0.5 ml mixture of pentamine sky blue (5% W/V) and ^{125}I -HSA (1 μl /animal). 10 minutes later 3 intradermal injections of FMLP (10 μg /site), 1 injection of histamine (1 μg /site) and 1 injection of vehicle (100 μl of 0.2% DMSO V/V in Hanks Buffered salt solution) were made on the left hand side of the animal (preinjection sites). 5 minutes later the drug (1 ml/kg) or the vehicle (50% PEG 400 V/V in distilled water, 1 ml/kg) was administered (i.v.). 10 minutes later an identical pattern of intradermal injections was made on the opposite flank of the animal (post-injection sites). These responses were allowed to develop for 15 minutes before the animal was sacrificed and a blood sample taken.

Skin sites and plasma samples were counted for 1 minute on a gamma counter and the degree of oedema calculated as μl plasma/skin site. Statistical analysis was done by a paired t-test on the mean of the 3 pre-injection site values of μl plasma obtained for FMLP/animal. The percentage inhibition of drug or vehicle was calculated as follows

$$X \% = \left(1 - \frac{\bar{X} \mu\text{l plasma (post-injection site)}}{\bar{X} \mu\text{l plasma (pre-injection site)}} \right) \times 100$$

Example No.	% inhibition	(mg/kg)
30	40.0	(1)

6. Effect of orally administered compounds on the FMLP-induced skin oedema of guinea-pigs *in vivo*-Test's p.o.

Guinea-pigs (600-800 g) were fasted overnight and orally treated with vehicle (1% Tylose w/v at 5 ml/kg) or drug (10 mg/kg; 2 mg/ml in 1% Tylose at 5 ml/kg) 40 minutes later the animals were anesthetized with pentobarbitone sodium (40 mg/kg; i.p.) and 0.6 ml of a mixture of pontamine sky blue (5% w/v) and ¹²⁵I-HSA (1 μ Ci/animal) was injected (i.v.). 90 minutes after oral pretreatment FMLP (50 μ g/site) was injected (i.d.) at 4 different sites, histamine (1 μ g/site) and vehicle (100 μ l, 1% DMSO v/v in Hanks buffered salt solution) were both injected (i.d.) at 2 different sites.

The responses were allowed to develop for 30 minutes before the animal was sacrificed and a blood sample taken.

Skin sites and plasma samples were counted for 1 minute on a gamma counter. The degree of oedema was calculated as μ l plasma/skin site.

Statistical analysis was carried out by a Mann-Whitney U-test on the mean of the 4 values of μ l Plasma obtained for FMLP/animal.

Example No.	% inhibition	(mg/kg)
30	46	(25)

The new active compounds can be converted in a known manner into the customary formulations, such as tablets, coated tablets, pills, granules, aerosols, syrups, emulsions, suspensions and solutions, using inert, nontoxic, pharmaceutically suitable excipients or solvents. In this connection, the therapeutically active compound should in each case be present in a concentration of about 0.5 to 90% by weight of the total mixture, i.e. in amounts which are sufficient in order to achieve the dosage range indicated.

The formulations are prepared, for example, by extending the active compounds with solvents and/or excipients, if appropriate using emulsifiers and/or dispersants, where, for example, in the case of the use of water as a diluent, organic solvents can be used as auxiliary solvents if appropriate.

Administration is carried out in a customary manner, preferably orally or parenterally, in particular perlingually or intravenously.

In the case of parenteral administration, solutions of the active compound can be employed using suitable liquid vehicles.

In general, it has proved advantageous on intravenous administration to administer amounts from about 0.001 to 10 mg/kg, preferably about 0.01 to 5 mg/kg of body weight to achieve effective results, and on oral administration the dosage is about 0.01 to 25 mg/kg, preferably 0.1 to 10 mg/kg of body weight.

In spite of this, it may be necessary to depart from the amounts mentioned, in particular depending on the body weight or the type of application route, on individual behaviour towards the medicament, the manner of its formulation and the time or interval at which administration takes place. Thus, in some cases it may be sufficient to manage with less than the abovementioned minimum amount, while in other cases the upper limit mentioned must be exceeded. In the case of administration of relatively large amounts, it is advisable to divide these into several individual doses over the course of the day.

Solvents

- I petrolether : ethylacetate 6:1
- II petrolether : ethylacetat 5:1
- III petrolether : ethylacetate 5:2


- IV dichloromethane : methanol 95:5
 V dichloromethane : methanol 9:1
 VI dichloromethane
 DMF dimethylformamide

5

Starting compoundsExample I

10 2'-Hydroxy-3-oxo-4'-[tetrahydro-2H-pyran-2-yl]oxy]benzenebutanoic acid methylester

15



20 20.0 g (0,089 mol) 2',4'-Dihydroxy-3-oxo-benzenebutanoic acid methylester were dissolved in 200 ml dichloromethane / tetrahydrofuran (95:5) and 9.2 ml (0.1 mol) 3,4-dihydro-2H-pyran and 10 mg p-toluenesulfonic acid were added successively. The suspension was stirred at room temperature for 1 hour. 400 ml of a NaHCO₃ solution were added, the organic layer separated and washed three times with water. The organic phase was dried using Na₂SO₄ and concentrated in vacuo. The residue was recrystallised from

25 diethylether.

Yield: 13.4 g (49% of theory)


R_f = 0,55, I

Example II

30

4-Benzyloxy-2-hydroxyacetophenone

35



40

Equivalent amounts, 152.15 (1.0 mol) of 2,4-dihydroxyacetophenone and 118.9 ml (1.0 mol) of benzylbromide were dissolved in 1.2 l acetone, 138 g potassium carbonate were added, and the mixture was stirred under reflux for 5 hours. Subsequently it was filtered off, the mother liquor was concentrated in vacuo and the residue recrystallised from diethylether.

45 Yield: 197 g (81%)


R_f = 0.82, III

50

55

Example III

6-Benzyloxy-3-methyl-2-(4-methylbenzoyl)benzo[b]furan




Equivalent amounts, 137.7 g (0,47 mol) 4-benzyloxy-2-hydroxy-acetophenone and 100 g (0,47 mol) ω -bromo-4-methylacetophenone were stirred under reflux for 12 h in 700 ml acetone in the presence of 65 g K_2CO_3 . The mixture was filtered off, the solvent removed in vacuo and the residue recrystallised from diethylether.

Yield: 94 g (56%)

$R_f = 0.63$, II

Example IV

6-Benzyloxy-3-bromo-methyl-2-(4-methylbenzoyl)benzo[b]furan



39.1 g (0.11 mol) of the compound from example III were dissolved in 500 ml carbon tetrachloride, 21.5 g (0.12 mol) of N-bromosuccinimide were added and the mixture was treated with 0.3 g dibenzoyl peroxide and heated to reflux for 12 hours. The mixture was filtered while hot, the solvent was distilled off in vacuo and the residue was purified by chromatography.

Yield: 19.4 g (41%)

$R_f = 0.8$, VI


Exempl V and VI

6-Benzyloxy-2-(4-methylbenzoyl)-3-benzofuran-carboxaldehyde (V)

5

10

15



10.4 g NaHCO_3 in 60 ml DMSO were heated under argon to 150°C . 7.0 g (16 mmol) of 6-benzyloxy-3-bromo-methyl-2-(4-methylbenzoyl)benzo[b]furan, dissolved in 60 ml DMSO, were added within 1 min. After 15 min at 150°C the mixture was poured onto ice and subsequently extracted three times with ethylacetate/diethylether (1:1). The organic phase was washed twice with H_2O , once with a NaCl solution and dried over MgSO_4 . The solvent was removed in vacuo and the residue purified by chromatography.


Yield: 5.4 g (91%)

$R_f = 0.74$, VI

By adding the benzylbromide (example IV) in a solid form the 6-benzyloxy-3-hydroxymethyl-2-(4-methylbenzoyl)benzo[b]furan (VI) was isolated in changing yields.

30

35




40 $R_f = 0.17$, II

Example VII

Methyl 2'-hydroxy-3-oxo-5'-[(tetrahydro-2H-pyran-2-yl)oxy]benzene butanoate

45

50




55 The compound has prepared in analogy to the procedure of Example I.

Yield: 56 %

$R_f = 0.58$.

Preparation ExamplesExample 1

5 2-(4-Bromo-benzoyl)-6-[(tetrahydro-2H-pyran-2-yl)oxy]-3-benzofuranpropanoic acid methylester




Equivalent amounts, 1.5 g (4.9 mmol) of 2'-Hydroxy-3-oxo-4'-[(tetrahydro-2H-pyran-2-yl)oxy]benzenebutanoic acid, methylester and 1.35 g (4.9 mmol) of ω -bromo-4-bromoacetophenone were dissolved in 50 ml acetone and 1.35 g (9.7 mmol) of potassium carbonate were added. The suspension was heated under reflux for 16 hours. The mixture was filtered, the solvent was distilled off in vacuo and the residue was taken up in ethylacetate. The organic phase was washed three times with water, one time with a NaCl solution, dried over Na₂SO₄ and concentrated in vacuo. The residue was further purified by chromatography (silica gel: Me₂SO 60).

Yield: 1.54 g (65.1%)

R_f = 0.53, I

The compounds shown in Tables 1 and 2 are prepared in analogy to the procedure of Example 1:

Table 1:




Ex.-No.	A	D	E	L	Solvent	R _f [*]	Yield (% of theory)
2	H	H	NO ₂	H		0.72, III	29
3	H	H	OCH ₃	H		0.48, II	50
4	H	H	Cl	H	DMF	0.45, II	85
5	H	H	F	H		0.63, II	48
6	H	H	CF ₃	H		0.65, II	20
7	H	H	CN	H	DMF	0.45, III	52
8	H	H	OCHF ₂	H	DMF	0.32, II	44
9	H	NO ₂	H	H	DMF	0.49, III	48

Table 1:

Ex.-No.	A	D	E	L	Solvent	R _f [*]	Yield (% of theory)
10	H	CN	H	H		0.48, III	30
11	H	CN	NH ₂	H		0.38, III	22
12	OCH ₃	H	H	OCH ₃		0.49, III	53
13	H	Cl	NHCOCH ₃	H		0.30, III	50
14	H	Br	H	H		0.65, III	53
15	H	H	CH ₃	H		0.67, II	55
16	H	Cl	H	H		0.70, III	46
17	H	H	COOCH ₃	H		0.56, III	60
18	H	Cl	NH ₂	Cl		0.53, III	38
19	Cl	H	Cl	H		0.50, II	75

Table 2:




Ex.-No.	A	D	E	a	Solvent	R _f [*]	Yield (% of theory)
20	H	H	CH ₃	2		0.57, VI	16
21	H	H	F	2		0.66, II	23
22	H	H	CH ₃	3		0.68, VI	quant.
23	H	NO ₂	H	3	MIBK	0.68, II	32
24	H	H	NO ₂	2	MIBK	0.41, II	7
25	H	NO ₂	H	2	MIBK	0.65, II	63
26	H	H	CN	2		0.39, III	62

MIBK = methylisobutylketone

Example 27

2-(4-Bromo-benzoyl)-6-hydroxy-3-benzofuranpropanoic acid methylester




4.2 g (8.64 mmol) of 2-(4-Bromo-benzoyl)-6-[(tetrahydro-2H-pyran-2-yl)oxy]-3-benzofuranpropanoic acid methylester were dissolved in 100 ml methanol and 10 mg p-toluene-sulfonic acid were added. The suspension was stirred at r.t. for 2 hours. The solvent was distilled off, the residue solved in ethylacetate and washed two times with water, once with a NaHCO₃ solution and once with a NaCl solution. The organic layer was dried using Na₂SO₄, concentrated in vacuo and the residue was further purified by chromatography (silica gel 60).

Yield: 3.0 g (86%)

R_f: 0.30, III

The compounds shown in Table 3 are prepared in analogy to the procedure of Example 27:

Table 3:



Ex.-No.	A	D	E	L	a	R _f [*]	Yield (% of theory)
28	H	H	NO ₂	H	2	0.27, III	80
29	H	H	OCH ₃	H	2	0.46, III	85
30	H	H	Cl	H	2	0.31, III	50
31	H	H	F	H	2	0.10, II	89
32	H	H	CF ₃	H	2	0.14, II	61
33	H	H	CN	H	2	0.73, V	94
34	H	H	OCHF ₂	H	2	0.08, II	96
35	H	NO ₂	H	H	2	0.21, III	70

Table 3:


Ex.-No.	A	D	E	L	a	R _f [*]	Yield (% of theory)
36	H	CN	H	H	2	0.10, II	70
37	H	CN	NH ₂	H	2	0.08, III	81
38	OCH ₃	H	H	OCH ₃	2	0.17, III	96
39	H	Cl	NHCOCH ₃	H	2	0.77, V	82
40	H	Br	H	H	2	0.37, III	65
41	H	H	CH ₃	H	2	0.09, II	63
42	H	Cl	H	H	2	0.43, III	10
43	H	H	COOCH ₃	H	2	0.24, III	37
44	H	Cl	NH ₂	Cl	2	0.32, III	87

Table 3:

Ex.-No.	A	D	E	L	a	R _f [*]	Yield (% of theory)
45	H	CH ₃	CH ₃	H	2	0.09, II	90
46	H	CH ₃	H	H	2	0.10, II	95
47	H	H	CH ₂ CH ₃	H	2	0.10, III	85
48	H	H	CH ₃	H	4	0.45, IV	90
49	H	Cl	Cl	H	2	0.20, III	89
50	Cl	H	Cl	H	2	0.35, III	45

Example 51

6-Hydroxy-2-(4-methyl-benzoyl)-3-benzofuranpropanoic acid



1.5 g (4.4 mmol) of the compound from example 41 were dissolved in 50 ml methanol/tetrahydrofuran (1:1) and 5.5 ml of a 2 N NaOH solution were added. The mixture was stirred at r.t. for 24 hours, dissolved in

water and acidified with 1 N hydrochloric acid. The precipitate was filtered off, washed several times with water and dried in vacuo.

Yield: 1.4 g (97%)

R_f: 0.29, V

- 5 The compounds shown in Table 4 are prepared in analogy to the procedure of example 51:

Ex.-No.	A	D	E	R ²	a	R _f [*]	Yield (% of theory)
52	H	CH ₃	H	-OH	2	0.27, V	quant.
53	H	CH ₃	CH ₃	-OH	2	0.13, IV	quant.
54	H	H	CH ₂ CH ₃	-OH	2	0.13, IV	quant.
55	H	H	CH ₃	-OCH ₂ -C ₆ H ₅	2	0.54, V	94
56	H	H	CH ₃	-Cl	2	0.37, V	57
57	H	H	OCH ₃	-OH	2	0.41, V	quant.
58	H	H	Cl	-OH	2	0.55, V	95
59	H	H	F	-OH	2	0.50, V	90

Table 4:





Table 4:

Ex.-No.	A	D	E	R ²	a	R _f [*]	Yield (% of theory)
60	H	H	Br	-OH	2	0.55, V	quant.
61	H	H	CF ₃	-OCH ₃	2	0.56, IV	94
62	H	H	CN	-OCH ₃	2	0.54, IV	72
63	H	H	Br	-OH	4	0.20, IV	95
64	H	H	CH ₃	-OH	4	0.58, V	94

Example 65

2-(4-Cyano-benzoyl)-6-methoxy-3-benzofuran-propanoic acid methylester




0.2 g (0.57 mol) of the compound from example 33 were dissolved in 10 ml acetone and subsequently 100 mg potassium carbonate and 0.054 ml (0.57 mmol) dimethylsulfate were added. The mixture was heated under reflux for 1 hour, the solvent removed in vacuo and the residue washed several times with diethylether. The product was further purified, if appropriate, by chromatography.

Yield: 0.13 g (63%)

$R_f = 0.60$, III

The compounds shown in Table 5 are prepared in analogy to the procedure of Example 65:

Table 5:



Ex.-No.	A	D	E	L	R ⁴	R _f [*] (% of theory)	Yield (% of theory)	Solvent	M.P. (°C)
66	H	H	Cl	H	CH ₃	0.70, III	95		
67	H	H	Br	H	CH ₃	0.69, III	quant.		
68	H	H	CF ₃	H	CH ₃	0.70, III	quant.		
69	H	CN	H	H	CH ₃	0.49, III	quant.		
70	OCH ₃	H	H	OCH ₃	CH ₃	0.35, III	94		
71	H	CN	NH ₂	H	CH ₃	0.18, III	71		
72	H	NO ₂	H	H	CH ₃	0.47, III	62		
73	H	Cl	H	H	CH ₃	0.65, III	quant.		
74	H	Cl	NH ₂	Cl	CH ₃	0.74, III	quant.		

Table 5:

Ex.-No.	A	D	E	L	R ⁴	R _f [*]	Yield (% of theory)	Solvent	M.P. (°C)
75	Cl	H	Cl	H	CH ₃	0.65, III	95		
76	H	H	CN	H	CH ₂ -CO-OCH ₃	0.23, III	91		
77	OCH ₃	H	H	OCH ₃	-CH ₂ -CO-OCH ₃	0.29, III	84	DMF	50
78	H	CN	NH ₂	H	-CH ₂ -CO-OCH ₃	0.09, III	56	DMF	50
79	H	Br	H	H	-CH ₂ -CO-OCH ₃	0.49, III	quant.	DMF	50
80	H	Br	H	H	-CH ₂ -CO-OC ₂ H ₅	0.59, III	85	DMF	50
81	H	Br	H	H	-CH ₂ -CO-CH ₃	0.44, III	92		
82	H	H	NO ₂	H	-CH ₂ -CO-OCH ₃	0.35, III	quant.		
83	H	H	NO ₂	H	-CH ₂ -CO-OC ₂ H ₅	0.48, III	quant.		
84	H	NO ₂	H	H	-CH ₂ -CO-OCH ₃	0.23, III	85		
85	H	NO ₂	H	H	-C ₅ H ₉	0.65, III	79		
86	H	Cl	H	H	-CH ₂ -CO-OCH ₃	0.39, III	88		

Table 5:


Ex.-No.	A	D	E	L	R ⁴	R _f [*]	Yield (% of theory)	Solvent	M.P. (°C)
87	H	Cl	H	H	-C ₅ H ₉	0.58, II	quant.		
88	H	H	CH ₃	H	-CH ₃	0.25, II	77		
89	H	H	CH ₃	H	-CH ₂ -CO-CH ₃		78		57
90	H	H	CH ₃	H	-CH ₂ -CO-C ₄ H ₉		41		53
91	H	H	CH ₃	H	-C ₅ H ₉		63		64
92	H	H	CH ₃	H	-CH ₂ -CO-OCH ₃	0.07, II	68		
93	H	H	CH ₃	H	-CH ₂ -C ₆ H ₅	0.60, VI	80		
94	H	H	OCH ₃	H	-CH ₂ -C ₆ H ₅	0.40, IV	85		
95	H	Cl	NH-CO-CH ₃	H	-CH ₂ -CO-OCH ₃	0.10, III	70		
96	H	Cl	Cl	H	-CH ₂ -CO-OCH ₃	0.20, II	89	DMF	
97	H	Cl	Cl	H	-CH ₂ -CO-OC ₂ H ₅	0.28, II	quant.	DMF	
98	H	H	CH ₃	H			81	DMF	98

Table 5:




Ex.-No.	A	D	E	L	R ⁴	Yield (% of theory)	Solvent	M.P. (°C)
99	H	H	CH ₃	H		61	DMF	126
100	H	H	CH ₃	H		59	DMF	102
101	H	H	CH ₃	H	-CH ₂ -CO-C ₂ H ₅	81	DMF	89
102	H	H	Cl	H	-CH ₂ -CO-OCH ₃	79	DMF	83
103	H	H	Cl	H	-C ₅ H ₉	58	DMF	48
104	H	H	Cl	H	-CH ₂ -CO-CH ₃	50	DMF	75
105	H	H	CH ₃	H	-C ₂ H ₅	86		65
106	H	H	Cl	H	-C ₂ H ₅	84	DMF	59
107	H	H	Cl	H		70	DMF	108

Table 5:



Ex.-No.	A	D	E	L	R ⁴	Yield (% of theory)	Solvent	M.P. (°C)	R _f [*]
108	H	H	CH ₃	H	-CH ₂ -CO-OC ₂ H ₅	62		115	
109	H	H	Cl	H	-CH ₂ -CO-OC ₂ H ₅	45	DMF	72	
110	H	H	Cl	H	-CH ₂ -CO-C ₂ H ₅	84	DMF	86	0.39, III
111	H	H	Cl	H	-CH ₂ -CO- 	30	DMF		0.44, III
112	H	H	Cl	H	-C ₃ H ₆ -COOCH ₃	56	DMF	56	0.43, III
113	H	H	Cl	H	-CH ₂ -C(=CH ₂)-COOCH ₃	81	DMF	96	0.28, II
114	H	H	Cl	H	-CH ₂ -C(=CH ₂)-COOH	57	DMF	145	0.29, IV
115	H	H	Cl	H	-CH ₂ -CH=CH-COOCH ₃	47	DMF	62	0.44, III

Table 5:

Ex.-No.	A	D	E	L	R ⁴	Yield (% of theory)	M.P. (°C)	Solvent	R _f [*]
116	H	H	Cl	H	-CH(CN)-CH ₃	45		DMF	0.55, III
117	H	H	Cl	H	-H ₂ C-C(=O)-NH ₂	60	166	DMF	0.44, IV

Example 118

6-Methoxy-2-(4-tetrazolyl-benzoyl)-3-benzofuran-propanoic acid methylester



0.95 g (2.62 mmol) of the compound from example 65 w r dissolved in 15 ml DMF, 0.85 g (13.1 mmol) sodium azide and 1.80 g (13.1 mmol) triethylamine hydrochloride wer added and the mixtur was heated under reflux for 24 h. After cooling at room temperature the mixtur was diluted with diethylether and subsequently washed three times with H₂SO₄ (1M), three times with water and 2 times with a NaCl solution.


5 The organic phase was dried using MgSO₄, the solvent was removed in vacuo and the residue purified by chromatography using dichloromethane/methanol (9:1).

Yield: 0.79 g (74%)

R_f: 0.09, V

10 Example 119


6-chloro-2-(4-tetrazolyl-benzoyl)-3-benzofuran-propanoic acid methylester



The title compound can be prepared in analogy to the procedure of Example 118.

30 Example 120

6-Hydroxy-2-(4-methyl-benzoyl)-3-benzofuran-propanoic acid ethylester



0.4 g (1.23 mmol) of the acid from example 51 were dissolved in 25 ml trichloromethane and 1.2 g p-toluene-sulfonic acid and 5 ml ethanol were added. The mixture was stirred under reflux for 24 h using a water separator. Subsequently the mixture was washed two times with water, dried over Na₂SO₄ and concentrated in vacuo.

Yield: quant.

R_f: 0.57, IV

50 The compounds shown in Table 6 are prepared in analogy to the procedure of Example 120:





Table 6:

Ex.-No.	A	D	E	R	R _f	Yield (% of theory)
121	H	CH ₃	CH ₃	-CH ₂ CH ₃	0.64, V	98
122	H	CH ₃	CH ₃	-CH(CH ₃) ₂	0.68, V	89
123	H	H	CH ₃	-C ₆ H ₉	0.56, IV	quant.

Example 124

6-Hydroxy-2-(4-methyl-benzoyl)-3-benzofuranpropanamide



0.5 g (1.54 mmol) of the acid from example 51 were dissolved in 5 ml THF, 1.06 g (6.55 mmol) 1,1'-carbonyl-bis-1H-imidazole were added and the mixture was stirred at room temperature for 12 hours. Subsequently NH₃-gas was added for 2 hours using an inlet pipe. After an additional hour stirring at r.t. the


solvent was distilled off in vacuo. The residue was taken up in ethylacetate and washed three times with water, one time with a NaHCO_3 solution and one time with a NaCl solution. The organic phase was dried using MgSO_4 and the solvent was removed in vacuo.

Yield: quant.

5 R_f : 0.42, V

The compounds shown in Table 7 are prepared in analogy to the procedure of Example 124:


Table 7:



Ex.-No.	R	R_f^*	Yield (% of theory)
125	-NH-CH ₃	0.38, V	97
126	-N(CH ₃) ₂	0.34, V	94

The compounds shown in Table 8 are prepared in analogy to the procedure of Example 65:


Table 8:



Ex.-No.	R ⁴	R_f^*	Yield (% of theory)
127	-CH ₃	0.21, II	30
128	-CH ₂ -C ₆ H ₅	0.28, II	40
129	-CH ₂ -C ₆ H ₄ -pNO ₂	0.13, II	68
130	-CH ₂ -C ₆ H ₄ -pCOOCH ₃	0.14, II	62
131	-C ₅ H ₉	0.34, II	56

Example 132


6-Benzyloxy-2-(4-methyl-benzoyl)-3-[3-oxo-3-(1-pyrrolidinyl)propenyl]benzo[b]furan



0.5 g (1.35 mmol) of the compound from example V were dissolved in 5 ml THF, cooled to -70°C and 0.8 ml (2.0 mmol) of *n*-BuLi (2.5 M solution in hexane) were added dropwise. Subsequently the mixture was stirred for 30 min. at -70°C and 0.50 g (2.0 mmol) of O,O-diethyl-[2-oxo-2-(1-pyrrolidinyl)ethyl]-phosphonacid ester, solved in 5 ml THF, were added dropwise. After stirring for 30 min. at -70°C the cooling bath was removed. After warming to 0°C 10 ml of a NH_4Cl solution were added. The mixture was extracted with ethylacetate and the organic phase was washed three times with water, one time with a NaCl solution and dried using MgSO_4 . The solvent was distilled off and the residue purified by chromatography. Yield: 0.45 g (72%)

R_f: 0.51, IV


The compounds shown in Table 9 are prepared in analogy to the procedure of Example 132:

Table 9:


Ex.-No.	E	R _f [*]	Yield (% of theory)
133	-COOCH ₃	0.19, III	60
134	-NO ₂	0.28, III	65

Example 135

Methyl 5-hydroxy-2-(4-methyl-benzoyl)-3-benzofuran-propionate



The title compound is prepared according to the process A starting from the compound of example VII.

Table 10

Ex. No.	A	D	E	L	X	Solvent	R _f	Yield (% of theory)
136	H	Cl	Cl	H	H	DMF	0.43,II	40
137	CH ₃	-NH-CO-CH ₃	CH ₃	H	H	DMF	0.48,IV	39
138	H	CH ₃	Cl	H	H	DMF	0.42,II	38
139	F	H	H	H	F	DMF	0.20,III	38
140	F	H	F	H	H	DMF	0.45,II	20
141	CH ₃	-NH-CO-CH ₃	Cl	H	H	DMF	0.60,IV	34

Table 10 (Continuation)




Ex. No.	A	D	E	L	X	Solvent	R _f	Yield (% of theory)
142	H	H	-SO ₂ -CH ₃	H	H	DMF	0.26,III	59
143	H	H	-SCH ₃	H	H	DMF	0.38,II	73
144	H	H	-O-SO ₂ -CH ₃	H	H	DMF	0.24,III	75
145	H	H		H	H	DMF	0.37,III	60
146	H	NO ₂	Cl	H	H	DMF	0.66,III	13
147	OCH ₃	H	H	H	OCH ₃	DMF	0.15,II	70

Table 11 (Continuation)

Ex. No.	A	D	E	L	X	R _f	Yield (% of theory)
157	H	H		H	H	0.02 V	17
158	F	H	H	H	F	0.13 II	96
159	OCH ₃	H	H	H	-OCH ₃	0.63 IV	90

The compounds shown in Table 12 were prepared in analogy to the procedure of Example 51

Table 12









Ex. No.	A	D	E	R ⁴	a	R _f	Yield (% of theory)
160	H	H	Cl	-CH ₂ -CO-NH ₂	2	0.07 (IV)	75
161	H	H	Cl	-CH ₂ COOH	2	0.1 (IV)	95
162	H	H	Cl	 -CH ₂ CON	2		
163	H	H	Cl	 -CH ₂ CON	2	0.33 (IV)	64

Table 12 (Continuation)

Ex. No.	A	D	E	R ⁴	a	R _f	Yield (% of theory)
164	H	H	Cl		2	0.35 (IV)	66
165	H	H	Cl	-CH ₂ CON(C ₂ H ₅) ₂	2	0.40 (IV)	55
166	H	H	Cl	-CH ₂ CON(CH ₃) ₂	2	0.41 (IV)	11
167	H	Cl	H	-CH ₂ CONH ₂	2	0.08 (IV)	86
168	H	H	Br	-CH ₂ CONH ₂	2	0.07 (IV)	93
169	H	H	Cl		2	0.2 (IV)	46
170	H	H	Cl	-CH ₂ CONHCH ₂ CH=CH ₂	2	0.1 (IV)	90
171	H	H	Cl	-CH ₂ CONHC ₃ H ₇	2	0.1 (IV)	88
172	H	H	Cl	-CH ₂ CN	2	0.15 (IV)	80
173	H	H	CH ₃	-CH ₂ CN	2	0.16 (IV)	90
174	H	H	Cl	-CH ₂ CH ₂ OH	2	0.1 (IV)	85
175	H	H	Cl		2	0.2 (IV)	89

The compounds shown in Table 13 and Table 14 are prepared in analogy to the procedure of example 65

Table 13



Ex. No.	A	D	E	L	X	R ⁴	R _T	Yield (% of theory)
176	H	Cl	NH ₂	Cl	H	-CH ₂ -CO-OCH ₃	0.48, III	quant.
177	H	Cl	NH ₂	Cl	H	-CH ₂ -CO-NH ₂	0.45, V	86
178	CH ₃	-NH-CO-CH ₃	CH ₃	H	H	-CH ₂ -CO-OCH ₃	0.58, V	quant.
179	CH ₃	-NH-CO-CH ₃	CH ₃	H	H	-CH ₂ -CO-NH ₂	0.54, V	84
180	H	CH ₃	Cl	H	H	-CH ₂ -CO-OCH ₃	0.58, III	quant.
181	H	CH ₃	Cl	H	H	-CH ₂ -CO-NH ₂	0.60, V	91
182	H	Cl	NH ₂	Cl	H	-CH ₃	0.71, III	quant.
183	F	H	H	H	F	-CH ₂ -CO-OCH ₃	0.36, III	95

Table 13 (Continuation)



Ex. No.	A	D	E	L	X	R ⁴	R _f	Yield (% of theory)
184	F	H	H	H	F	-CH ₂ -CO-NH ₂	0.65, V	98
185	F	H	F	H	H	-CH ₂ -CO-OCH ₃	0.19, III	97
186	F	H	F	H	H	-CH ₂ -CO-NH ₂	0.53, IV	92
187	CH ₃	-NH-CO-CH ₃	Cl	H	H	-CH ₂ -CO-OCH ₃	0.56, IV	75
188	CH ₃	-NH-CO-CH ₃	Cl	H	H	-CH ₂ -CO-NH ₂	0.28, IV	72
189	H	H	-S-CH ₃	H	H	-CH ₂ -CO-OCH ₃	0.17, III	87
190	H	H	-S-CH ₃	H	H	-CH ₂ -CO-NH ₂	0.53, IV	quant.
191	H	H	-SO ₂ -CH ₃	H	H	-CH ₂ -CO-OCH ₃	0.9, IV	86
192	H	H	-SO ₂ -CH ₃	H	H	-CH ₂ -CO-NH ₂	0.32, IV	67
193	H	H	-O-SO ₂ -CH ₃	H	H	-CH ₂ -CO-OCH ₃	0.79, IV	88
194	H	H	-O-SO ₂ -CH ₃	H	H	-CH ₂ -CO-NH ₂	0.42, IV	88
195	H	H	-SO ₂ -N ₂ O ₂	H	H	-CH ₂ -CO-OCH ₃	0.89, IV	88
196	H	H	-SO ₂ -N ₂ O ₂	H	H	-CH ₂ -CO-NH ₂	0.46, IV	89
197	H	-NO ₂	Cl	H	H	-CH ₂ -CO-OCH ₃	0.15, III	69
198	H	H		H	H	-CH ₂ -CO-NH ₂	0.42, V	quant.

Table 13 (Continuation)

Ex. No.	A	D	E	L	X	R ⁴	R _T	Yield (% of theory)
199	H	-CN	H	H	H	-CH ₂ -CO-OCH ₃	0.22, III	quant.
200	-Cl	H	Cl	H	H	-CH ₂ -CO-OCH ₃	0.25, II	95
201	H	H		H	H	-CH ₂ -CO-OCH ₃	0.75, V	35

5

10

15

20

25

30

35


40

45

50

55

Table 14



Ex. No.	A	D	E	L	R^4	R_f	Yield (% of theory)	m.p. °C
202	H	H	Cl	H	$—CH_2—C \equiv N$	0.5 (III)	76	75
203	H	H	Cl	H	$—(CH_2)_3—C \equiv N$	0.32 (III)	85	77
204	H	H	CH_3	H	$—CH_2—C(=O)—NH_2$	0.4 (IV)	87	160
205	H	H	NO_2	H	$—CH_2—C(=O)—NH_2$	0.4 (IV)	97	181
206	H	H	CH_3	H	$—CH_2—C(=O)—CH_2—COOH$	0.6 (IV)	66	123
207	H	NO_2	H	H	$—CH_2—C(=O)—NH_2$	0.15 (IV)	88	181

Table 14 (Continuation)

Ex. No.	A	D	E	L	R ⁴	R _T	Yield (% of theory)	m.p. °C
208	H	Br	H	H	$-\text{CH}_2-\text{C}(=\text{O})-\text{NH}_2$	0.24 (IV)	81	174
209	H	CN	H	H	$-\text{CH}_2-\text{C}(=\text{O})-\text{NH}_2$	0.17 (IV)	87	203
210	Cl	H	Cl	H	$-\text{CH}_2-\text{C}(=\text{O})-\text{NH}_2$	0.46 (IV)	48	161
211	H	H	CN	H	$-\text{CH}_2-\text{C}(=\text{O})-\text{NH}_2$	0.47 (IV)	85	206
212	H	H	Cl	H	$-\text{CH}_2-\text{COOH}$	0.52 (V)	36	
213	H	H	CH ₃	H	$-\text{CH}_2-\text{C}\equiv\text{N}$	0.37 (III)	96	
214	H	H	Cl	H	$-\text{CH}_2-\text{C}(\text{N}(\text{CH}_3))_2$	0.46 (IV)	quant.	
215	H	H	Cl	H	$-\text{SO}_2\text{CF}_3$	0.79 (III)	96	
216	H	H	CH ₃	H	$-\text{SO}_2\text{CF}_3$	0.75 (III)	82	

Table 14 (Continuation)







Ex. No.	A	D	E	L	R ⁴	R _f	Yield (% of theory)	m.p. °C
217	H	H	CH ₃	H		0.44 (IV)	90	
218	H	H	Cl	H	-CF ₂ H	0.34 (II)	48	
219	H	H	Cl	H	-CH ₂ CF ₃	0.44 (II)	20	96
220	H	H	Cl	H		0.4 (IV)	89	
221	H	H	Cl	H		0.23 (IV)	79	
222	H	H	Cl	H		0.31 (IV)	89	
223	H	H	Cl	H		0.43 (IV)	94	123-124
224	H	H	Cl	H		0.71 (IV)	96	

Table 14 (Continuation)

Ex. No.	A	D	E	L	R ⁴	R _f	Yield (% of theory)	m.p. °C
225	H	Cl	H	H	-CH ₂ -CO-NH ₂	0.37 (IV)	78	185
226	H	H	Br	H	-CH ₂ -CO-NH ₂	0.46(IV)	72	160
227	H	H	Cl	H	$\begin{array}{c} \text{O} \quad \text{O} \\ \diagdown \quad \diagup \\ \text{---CH}_2\text{---C---} \end{array}$	0.15 (VI)	30	114
228	H	H	Cl	H	-CH ₂ CH ₂ Cl	0.79 (III)	16	224
229	H	H	Cl	H	-CH ₂ CH ₂ OH	0.21 (III)	32	96
230	H	H	Cl	H	$\begin{array}{c} \text{O} \\ \parallel \\ \text{---CH}_2\text{---C---N---} \end{array} \begin{array}{l} \text{CH}_3 \\ (\text{CH}_2)_3\text{CH}_3 \end{array}$	0.8 (IV)	90	
231	H	H	Cl	H	$\begin{array}{c} \text{O} \\ \parallel \\ \text{---CH}_2\text{---C---N---} \end{array} \begin{array}{l} \text{C}_2\text{H}_5 \\ (\text{CH}_2)_3\text{CH}_3 \end{array}$	0.8 (IV)	96	

Table 14 (Continuation)








Ex. No.	A	D	E	L	R ⁴	R _f	Yield (% of theory)	m.p. °C
232	H	H	Cl	H		0.56 (IV)	85	123-4
233	H	H	Cl	H		0.74 (IV)	93	
234	H	H	Cl	H		0.84 (IV)	10	
235	H	H	Cl	H		0.57 (IV)	95	144-146
236	H	H	Cl	H		0.84 (IV)	64	126-129
237	H	H	Cl	H		0.91 (IV)	96	99-101

Table 14 (Continuation)


Ex. No.	A	D	E	L	R ⁴	R _f	Yield (% of theory)	m.p. °C
238	H	H	Cl	H	$\text{---CH}_2\text{---C(=O)---N(C}_4\text{H}_9)_2$	0.91 (IV)	91	
239	H	H	Cl	H	$\text{---CH}_2\text{---C(=O)---N(H)(i-Bu)}$	0.9 (IV)	72	133-135
240	H	H	Cl	H	$\text{---CH}_2\text{---C(=O)---N(H)CH}_3$	0.52 (IV)	30	135-137
241	H	H	Cl	H	$\text{---CH}_2\text{---C(=O)---N(H)CH}_2\text{---CH=CH}_2$	0.73 (IV)	75	129
242	H	H	Cl	H	$\text{---CH}_2\text{---C(=O)---N(H)C}_3\text{H}_7$	0.64 (IV)	61	121
243	H	H	Cl	H	-COCH ₃	0.27 (II)	67	

Table 14 (Continuation)

Ex. No.	A	D	E	L	R ⁴	R _f	Yield (% of theory)	m.p. °C
244	H	H	Cl	H	-CH ₂ CONHCH ₂ CO ₂ C ₂ H ₅	0.4 (V)	62	114-116
245	H	H	CH ₃	H		0.8 (III)	10	
246	H	H	Cl	H	CH ₂ CONHCH ₂ OH	0.4 (III)	22	128

The compounds shown in Table 15 are prepared in analogy to the procedure of Example 65


Table 15



Ex. No.	R ⁴	R _f	Yield (% of theory)
247	-CH ₂ -CO-OCH ₃	0.71, IV	74
248	-CH ₂ -CO-OC ₂ H ₅	0.72, IV	78
249	-CH ₂ -CO-NH ₂	0.30, IV	20

The compounds shown in Table 16 are prepared in analogy to the procedure of Example 132


Table 16:



Ex. No.	R ₄	E	R _f	Yield (% of theory)
250	H ₃ C-	Cl	0.42, II	61

Exempl 251

2-(4-Chloro-benzoyl)-6-methoxy-3-benzofuranpropionitril




0.49 g (1.45 mmol) of the compound from example 250 was suspended in 10 ml methanol and hydrogenated for 1 hour at 3.5 bar and room temperature in the presence of 60 mg palladium-on-charcoal catalyst (5%). The catalyst was filtered off and the residue was evaporated. The product was further purified, if appropriate, by chromatography.

Yield: 71%

R_f = 0.40, II

Example 252


2-(4-Chloro-benzoyl)-6-hydroxy-3-benzofuranpropionitril



The title compound is prepared according to example 251.

Example 253

(4-Chloro-phenyl)-{6-methoxy-3-[2-(2H-tetrazol-5-yl)-ethyl]-benzofuran-2-yl}-methanone




0.2 g (0.59 mmol) of the compound from example 251 were dissolved in 5 ml xylene and 0.17 ml (0.59 mmol) tributyltin chloride and 38.4 mg sodium azide were added. The resulting mixture was heated at 80 °C under argon atmosphere for 3 days. After the mixture was cooled to ambient temperature, it was added with stirring to 10 ml of ice-cold, dry methanol saturated with HCl gas. The mixture was stirred for 90 min before it was concentrated in vacuo. The residue was purified by chromatography.

Yield: 49%

 R_f : 0.02, V

Example 254

3-[2-(4-Chloro-benzoyl)-6-[(tetrahydro-2H-pyran-2-yl)oxy]-benzofuran-3-yl]-N-(2-hydroxy-1,1-dimethyl-ethyl)-propionamide




3.0 g (7.0 mmol) of the acid from example 4 were dissolved in a mixture of 10 ml acetonitrile/10 ml pyridine and 623 mg (7.0 mmol) 2-amino-2-methylpropanol, 2.9 ml (21.0 mmol) triethylamine and 2.03 ml (21.0 mmol) tetrachloromethane were added. 5.5 g (21.0 mmol) triphenylphosphin dissolved in a mixture of 10 ml acetonitrile/10 ml pyridine were added dropwise. After stirring at room temperature for 12 h the mixture was diluted with water and extracted 3 x with ethylacetate. The organic phase was washed with a NaCl solution, dried using $MgSO_4$ and the solvent was removed in vacuo. The residue was purified by chromatography using dichloromethane/methanol (9:1).

Yield: 2.42 g (72%)

 R_f : 0.64, V.

Example 255

(4-Chloro-phenyl)-{3-[2-(4,4-dimethyl-4,5-dihydro-oxazol-2-yl)-ethyl]-6-hydroxybenzofuran-2-yl}-methanon




0.58 ml (8 mmol) thionylchloride were added dropwise under stirring to 1.0 g (2.0 mmol) of the compound from example 254. The mixture was stirred at room temperature for 12 hours and the excess thionylchloride was removed in vacuo. The residue was taken up in water and the pH of the solution was adjusted to pH = 8 by adding a 1 N sodium hydroxide solution. After extraction of the water phase 3 x with ethylether and removing of the solvent in vacuo the residue was purified by column chromatography.

Yield: 0.7 g (85%)

R_f: 0.70, V.

Example 256


2-{2-(4-Chloro-benzoyl)-3-[2-(4,4-dimethyl-4,5-dihydro-oxazol-2-yl)-ethyl]-benzofuran-6-yloxy}-acetamide





The title compound is prepared according to example 65.

The compounds shown in Table 17 are prepared by reaction of the carbon acids with NaOH.


Table 17



Ex. No.	R ⁴	A	D	E	R _f	Yield (% of theory)
257	-CH ₂ -CO-NH ₂	H	H	Cl	0.1 (V)	95
258	-CH ₂ -CO-N 	H	H	Cl	0.2 (V)	98
259	-CH ₂ -CO-N 	H	H	Cl	0.2 (V)	97
260	-CH ₂ -CO-N(C ₂ H ₅) ₂	H	H	Cl	0.15 (V)	98






The compounds shown in Table 18 are prepared in analogy to the procedure of example 65.

Table 18:




Ex. No.	A	D	E	R ⁴	R _f	Yield (% of theory)
261	H	H	CH ₃		0.58 (IV)	57
262	H	H	CH ₃		0.25 (IV)	80
263	H	H	CH ₃		0.36 (IV)	86
264	H	H	CH ₃		0.30 (IV)	74
265	H	H	CH ₃		0.33 (IV)	74

Table 18 (Continuation)

Ex. No.	A	D	E	R ⁴	R _f	Yield (% of theory)
266	H	H	Br		0.5 (IV)	86
267	H	H	Br		0.28 (IV)	75
268	H	H	Br		0.28 (IV)	86
269	H	H	Br		0.29 (IV)	65
270	H	H	Br		0.28 (IV)	80

The compounds shown in Table 19 were prepared in analogy to the procedure of Example 51.

Table 19:





Ex. No.	A	D	E	R ⁴	a	R _f	Yield (% of theory)
271	H	Cl	H	-OH	2	0.2 (IV)	83
272	H	H	Cl	$\text{---CH}_2\text{---C(=O)---N---CH}_2\text{---C}_6\text{H}_5$	2	0.15 (IV)	95
273	H	H	Cl	$\text{---CH}_2\text{---C(=O)---N(CH}_3\text{)---C}_4\text{H}_9$	2	0.3 (IV)	40
274	H	H	Cl	$\text{---CH}_2\text{---CH}_2\text{---N---}$ 	2	0.6 (V)	42
275	H	H	Cl	$\text{---CH}_2\text{---CH}_2\text{---N---}$ 	2	0.05 (V)	20

Table 19 (Continuation)






Ex. No.	A	D	E	R ⁴	a	R _T	Yield (% of theory)
276	H	H	Cl		2	0.5 (IV)	80
277	H	H	Br		2	0.42 (V)	50
278	H	H	Br		2	0.23 (V)	61
279	H	H	Br		2	0.29 (V)	79
280	H	H	Cl		2	0.25 (V)	67

Table 12 (Continuation)








Ex. No.	A	D	E	R ⁴	a	R _f	Yield (% of theory)
281	H	H	Br		2	0.27 (V)	85
282	H	H	CH ₃		2	0.54 (V)	58
283	H	H	CH ₃		2	0.26 (V)	88
284	H	H	CH ₃		2	0.44 (V)	47

Table 19 (Continuation)

Ex. No.	A	D	E	R ⁴	a	R _f	Yield (% of theory)
285	H	H	CH ₃		2	0.26 (V)	85
286	H	H	CH ₃		2	0.22 (V)	74


The compounds shown in Table 20 were prepared in analogy to the procedure of Example 120.



Ex. No.	A	D	E	R ¹⁰	R _I	Yield (% of theory)
287	H	H	Br	-O-C(CH ₃) ₃	0.2 (IV)	40
288	H	H	Br	-O-CH(CH ₃) ₂	0.22 (IV)	43

Example 289


2-[2-(4-Chloro-benzoyl)-3-(2-cyano-ethyl)-benzofuran-6-yl-oxy]-acetamide



The title compound is prepared according to example 65.

Example 290

3-[2-(4-Chloro-benzoyl)-6-hydroxy-benzofuran-3-yl]-propiohydroxamic acid




0.9 g (2.6 mmol) of the acid from example 58 were dissolved in THF. 1.85 g (11.1 mmol) carbonyl diimidazole were added and the reaction mixture was stirred at room temperature for 12 hours. 0.36 g (5.22 mmol) hydroxylamine hydrochloride were added and the mixture was stirred further 6 hours. The solvent was removed in vacuo and the residue was solved in ethylacetate. The organic phase was washed three times with water and one time with a NaHCO₃-solution and with a NaCl-solution. The organic phase was dried with Na₂SO₄ and concentrated in vacuo. The residue was further purified by chromatography.

Yield: 140 mg (15 %)

R_f: 0.29, V

Claims

1. Benzofuranyl- and -thiophenyl-alkane-carboxylic acid derivatives of the general formula



in which

R¹ and R²

are identical or different and represent hydrogen, halogen, carboxyl, cyano, nitro, trifluoromethyl or a group of a formula -OR⁴, -SR⁵ or -NR⁶R⁷,

in which

R⁶


denotes hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

R⁴, R⁵ and R⁷

are identical or different and

denote hydrogen, cycloalkyl having 3 to 6 carbon atoms, hydrogen or a 5 to 7-membered saturated or unsaturated heterocycle having up to 3 heteroatoms from the series comprising N, S and O, which are optionally substituted by identical or different substituents from the series comprising halogen, cyano, nitro or by straight-chain or branched alkyl or alkoxy-carbonyl each having up to 6 carbon atoms or


denote a residue of formula



or

denote straight-chain or branched alkyl or alkenyl each having up to 8 carbon atoms, and each of which is optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising trifluoromethyl, difluoromethyl, halogen, cyano, carboxy, hydroxy, straight-chain or branched alkoxy, alkoxycarbonyl or acyl each having up to 6 carbon atoms or by a 5-to 7-membered saturated or unsaturated heterocycle having up to 3 hetero atoms from the series comprising N, S and O and to which an aromatic ring can be fused,

or by N-methyl-substituted imidazolyl,
or by a residue of formula



or by phenyl,

wherein all rings are optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising nitro, halogen, carboxy or straight-chain or branched alkyl or alkoxycarbonyl each having up to 6 carbon atoms,

or alkyl or alkenyl are substituted by a group of formula -CO-NR⁸R⁹

in which

R⁸ and R⁹

are identical or different and denote, phenyl, adamantyl, cycloalkyl having up to 7 carbon atoms, benzyl, hydrogen, formyl, straight-chain or branched alkyl or alkenyl each having up to 6 carbon atoms and which are optionally substituted by carboxy, hydroxy or straight-chain or branched alkoxycarbonyl or up to 6 carbon atoms,

R⁸ and R⁹

together with the nitrogen atom form a 5 to 7 membered saturated or unsaturated heterocycle,

or

R⁴

denotes a protecting group of a hydroxyl group, difluoromethyl or a group of a formula -SO₂-X,

in which

X

denotes trifluoromethyl, phenyl or straight-chain or branched alkyl having up to 6 carbon atoms,

T


represents an oxygen or sulfur atom,

V

represents a straight-chain or branched alkylene or alkenylene chain each having 2 to 8 carbon atoms,

W

represents cyano, tetrazolyl or a group of a formula -CO-R¹⁰, -CO-NR¹¹R¹², -CONR¹³-SO₂-R¹⁴ or PO(OR¹⁵)(OR¹⁶), or a residue of the formula



5

10

15

20

25

30

35

40


45

- in which
- R^{10} denotes hydroxyl, cycloalkyloxy having up to 3 to 7 carbon atoms or straight-chain or branched alkoxy having up to 8 carbon atoms,
- R^{11} , R^{12} and R^{13} are identical or different and denote hydrogen, phenyl, or straight-chain or branched alkyl or acyl each having up to 6 carbon atoms and which are optionally substituted by hydroxyl,
- or
- R^{11} denotes hydrogen
- and
- R^{12} denotes hydroxyl,
- or
- R^{11} and R^{12} together with the nitrogen atom form a 5- or 6-membered saturated heterocycle,
- R^{14} denotes straight-chain or branched alkyl having up to 6 carbon atoms, which is optionally substituted by phenyl or trifluoromethyl, or
- denotes phenyl, which is optionally substituted by substituents from the series comprising halogen, cyano, nitro or by straight-chain or branched alkyl having up to 6 carbon atoms,
- R^{15} and R^{16} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms,
- R^3 represents phenyl, which is monosubstituted to trisubstituted by identical or different substituents from the series comprising hydroxyl, halogen, nitro, tetrazolyl, trifluoromethoxy, difluoromethoxy, trifluoromethyl, difluoromethyl, cyano, carboxy, straight-chain or branched alkyl, alkylthio, alkoxy, alkoxycarbonyl or acyl each having up to 8 carbon atoms or by a group of formula $-NR^{17}R^{18}$, $-(O)_aSO_2-R^{19}$ or $-SO_2NR^{20}R^{21}$
- in which
- a denotes a number 0 or 1,
- R^{17} and R^{18} have the meaning shown above for R^{11} and R^{12} and are identical to the latter or different from the latter,
- or
- R^{17} denotes hydrogen,
- and
- R^{18} denotes straight-chain or branched acyl having up to 6 carbon atoms
- and
- R^{19} has the abovementioned meaning of R^{14} and is identical to the latter or different from the latter,
- R^{20} and R^{21} have the above mentioned meaning of R^{11} and R^{12} and are identical to the latter or different from the latter.

2. Benzofuranyl- and -thiophenyl-alkane-carboxylic acid derivatives according to claim 1, wherein

- R^1 and R^2 are identical or different and represent hydrogen, fluorine, chlorine, bromine, nitro, trifluoromethyl or a group of a formula $-OR^4$, $-SR^5$ or $-NR^6R^7$,
- in which
- R^6 denotes hydrogen or straight-chain or branched alkyl having up to 3 carbon atoms,
- R^4 , R^5 and R^7 are identical or different and denote hydrogen, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, chinolyl, pyridyl, imidazolyl, 1,3-thiazolyl or thienyl, which are optionally substituted by identical or different substituents from the series comprising fluorine, chlorine, bromine, iodine, cyano, nitro or by straight-chain or branched alkyl or alkoxycarbonyl each having up to 5 carbon atoms or


d note a residue of formula



or

denote straight-chain or branched alkyl or alkenylen each having up to 6 carbon atoms, and each of which is optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising trifluoromethyl, difluoromethyl, fluorine, chlorine, bromine, iodine, cyano, carboxy, hydroxy, straight-chain or branched alkoxy, alkoxycarbonyl or acyl each having up to 5 carbon atoms or by chinolyl, pyridyl, pyrazolyl, 1,3-thiadiazolyl or thienyl, imidazolyl or N-methyl-substituted imidazolyl, and to which an aromatic ring can be fused,

or by a residue of formula



or by phenyl,

where in all rings are optionally monosubstituted to disubstituted by identical or different substituents from the series comprising nitro, fluorine, chlorine, bromine, iodine, carboxy or straight-chain or branched alkyl or alkoxycarbonyl each having up to 5 carbon atoms,

or alkyl or alkenylen are substituted by a group of formula $-CO-NR^8R^9$,

in which

R^8 and R^9 are identical or different and denote phenyl, adamantyl, cyclopropyl, cyclopentyl, benzyl, formyl, hydrogen, straight-chain or branched alkyl or alkenyl each having up to 5 carbon atoms and which are optionally substituted by carboxy, hydroxy or straight-chain or branched alkoxycarbonyl up to 4 carbon atoms or together with the nitrogen atom form a morpholinyl, piperidinyl, piperazinyl or pyrrolidinyl ring,

or

R^4 denotes acetyl, benzyl, tetrahydrofuranyl, difluoromethyl or a group of a formula $-SO_2-X$,


in which

X denotes trifluoromethyl, phenyl or methyl,

T represents an oxygen or sulfur atom,

V represents a straight-chain or branched alkylene or alkenylene chain each having 2 to 6 carbon atoms,


W represents cyano, tetrazolyl or a group of a formula $-CO-R^{10}$, $-CO-NR^{11}R^{12}$, $-CONR^{13}-SO_2R^{14}$ or $PO(OR^{15})(OR^{16})$, or a residue of the formula



- in which
- R^{10} denotes hydroxyl, cyclopropyloxy, cyclopentyloxy, cyclohexyloxy or straight-chain or branched alkoxy having up to 6 carbon atoms,
- R^{11} , R^{12} and R^{13} are identical or different and denote hydrogen, phenyl, straight-chain or branched alkyl or acyl each having up to 4 carbon atoms and which are optionally substituted by hydroxyl,
- or
- R^{11} denotes hydrogen
- and
- R^{12} denotes hydroxyl,
- or
- R^{11} and R^{12} together with the nitrogen atom form a pyrrolidinyl piperidinyl or a morpholinyl ring,
- R^{14} denotes straight-chain or branched alkyl having up to 5 carbon atoms, which is optionally substituted by phenyl or trifluoromethyl, or denotes phenyl, which is optionally substituted by substituents from the series comprising fluorine, chlorine, bromine, iodine, cyano, nitro or by straight-chain or branched alkyl having up to 4 carbon atoms,
- R^{15} and R^{16} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms,
- R^3 represents phenyl, which is monosubstituted to trisubstituted by identical or different substituents from the series comprising hydroxyl, fluorine, chlorine, bromine, iodine, nitro, trifluoromethoxy, trifluoromethyl, difluoromethoxy, difluoromethyl, cyano, carboxy, tetrazolyl, straight-chain or branched alkyl, alkylthio, alkoxy, alkoxycarbonyl or acyl each having up to 6 carbon atoms or by a group of formula $-NR^{17}R^{18}$, $-(O)_a-SO_2-R^{19}$ or $-SO_2-NR^{20}R^{21}$,
- in which
- a denotes a number 0 or 1,
- R^{17} and R^{18} have the meaning shown above for R^{11} and R^{12} and are identical to the latter or different from the latter,
- or
- R^{17} denotes hydrogen,
- and
- R^{18} denotes straight-chain or branched acyl having up to 6 carbon atoms
- and
- R^{19} has the abovementioned meaning of R^{14} and is identical to the latter or different from the latter,
- R^{20} and R^{21} have the above mentioned meaning of R^{11} and R^{12} and are identical to the latter or different from the latter.


3. Benzofuranyl- and -thiophenyl-alkane-carboxylic acid derivatives according to claim 1, wherein

- R^1 denotes hydrogen,
- R^2 denotes fluorine, chlorine, bromine, nitro, trifluoromethyl or a group of a formula $-OR^4$ or $-NR^6R^7$,
- in which
- R^4 denotes a group of a formula $-SO_2X$,
- in which
- X denotes phenyl, trifluoromethyl or methyl,
- or
- R^4 denotes hydrogen, difluoromethyl, tetrahydropyranyl, benzyl, acetyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, chinolyl, pyridyl, imidazolyl or thienyl, which are optionally substituted by identical or different substituents from the series comprising fluorine, chlorine, bromine, cyano, nitro or by straight-chain or branched alkyl or alkoxycarbonyl each having up to 4 carbon atoms or
- denotes a residue of the formula



or

denotes straight-chain or branched alkyl or alkenylene each having up to 5 carbon atoms, and each of which is optionally monosubstituted to disubstituted by identical or different substituents from the series comprising trifluoromethyl, difluoromethyl, fluorine, chlorine, bromine, cyano, carboxy, hydroxy, straight-chain or branched alkoxy, alkoxycarbonyl or acyl each having up to 5 carbon atoms or by chinolyl, imidazolyl or N-methyl substituted imidazolyl, pyridyl, and to which an aromatic ring can be fused, or by a residue of formula



or by phenyl,

wherein all rings are optionally monosubstituted to disubstituted by identical or different substituents from the series comprising nitro, fluorine, chlorine, bromine, carboxy or straight-chain or branched alkyl or alkoxycarbonyl each having up to 4 carbon atoms,

or alkyl or alkenylene are substituted by a group of formula $-CO-NR^8R^9$ in which

R^8 and R^9 are identical or different and denote phenyl, benzyl, adamantyl, cyclopropyl, cyclopentyl, formyl, hydrogen, straight-chain or branched alkyl or alkenyl each having up to 5 carbon atoms, which are optionally, substituted by carboxy, hydroxy or straight-chain or branched alkoxycarbonyl up to 3 carbon atoms

or

R^8 and R^9 together with the nitrogen atom form a morpholinyl, piperidinyl, piperazinyl or a pyrrolidinyl ring


R^6 denotes hydrogen, methyl or ethyl,

R^7 denotes hydrogen, methyl or ethyl,

T represents an oxygen or sulfur atom,

V represents a straight-chain or branched alkylene or alkenylene chain each having 2 to 5 carbon atoms,

W represents cyano, tetrazolyl, or a group of a formula $-CO-R^{10}$, $-CO-NR^{11}R^{12}$, $-CONR^{13}-SO_2-R^{14}$ or $PO(OR^{15})(OR^{16})$, or a residue of the formula



in which

R^{10} denotes hydroxyl, cyclopropyloxy, cyclopentyloxy, cyclohexyloxy or straight-chain or branched alkoxy having up to 5 carbon atoms,

R^{11} , R^{12} and R^{13} are identical or different and denote phenyl, hydrogen, straight-chain or

branched alkyl or acyl each having up to 4 carbon atoms and which are optionally substituted by hydroxyl,

or

R^{11} denotes hydrogen

and

R^{12} denotes hydroxyl,

R^{11} and R^{12} together with the nitrogen atom form a pyrrolidinyl piperidinyl or morpholinyl ring,

R^{14} denotes straight-chain or branched alkyl having up to 4 carbon atoms, which is optionally substituted by phenyl or trifluoromethyl, or denotes phenyl, which is optionally substituted by substituents from the series comprising fluorine, chlorine, bromine, cyano, nitro or by a straight-chain or branched alkyl having up to 3 carbon atoms,

R^{15} and R^{16} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 5 carbon atoms,

R^3 represents phenyl, which is monosubstituted to trisubstituted by identical or different substituents from the series comprising hydroxyl, fluorine, chlorine, bromine, nitro, tetrazolyl, trifluoromethyl, trifluoromethoxy, difluoromethoxy, difluoromethyl, cyano, carboxy, straight-chain or branched alkyl, alkylthio, alkoxy, alkoxycarbonyl or acyl each having up to 5 carbon atoms or by a group of formula $-NR^{17}R^{18}$, $-(O)_n-SO_2-R^{19}$ or $-SO_2-NR^{20}R^{21}$

in which

R^{17} and R^{18} have the meaning shown above for R^{11} and R^{12} and are identical to the latter or different from the latter,

or

R^{17} denotes hydrogen,

and

R^{18} denotes straight-chain or branched acyl having up to 5 carbon atoms

and


R^{19} has the abovementioned meaning of R^{14} and is identical to the latter or different from the latter,

R^{20} and R^{21} have the above mentioned meaning of R^{11} and R^{12} and are identical to the latter or different from the latter.

4. Benzofuranyl- and -thiophenyl-alkane-carboxylic acid derivatives according to claims 1 to 3 for controlling diseases.

5. Process for the preparation of benzofuranyl- and -thiophenyl-alkane-carboxylic acid derivatives to claims 1 to 3, characterized in that

[A] compounds of the general formula (II)



in which

R^1 , T , W and V have the above mentioned meaning,


R^{22} represents a group of formula $-OR^4$

in which


R^4 has the above mentioned meaning of R^4 , but does not represent hydrogen, are reacted with compounds of the general formula (III)

$R^3-CO-CH_2-Y$ (III)


in which
 R^3 has the above mentioned meaning,
 and
 Y represents a typical leaving group such as, for example, chlorine, bromine, iodine, tosylate or mesylate, preferably bromine, in inert solvents and in the presence of a base under cyclisation by customary methods
 [B] in the case, in which V represents alkenyl, compounds of the general formula (IV)




in which
 R^1 , R^3 , T and R^{22} have the above mentioned meaning,
 first are converted by reaction with N-bromosuccinimide, in inert solvents and in the presence of a catalyst to the compounds of the general formula (V)



in which
 R^1 , R^3 , T and R^{16} have the above mentioned meaning,
 and then by subsequent hydrolysis to compounds of the general formula (VI)



in which
 R^1 , R^3 , T and R^{22} have the above mentioned meaning,
 with in a last step are reacted with compounds of the general formula (VII)



in which
 R^{11} and R^{12} have the above mentioned meaning,
 R^{23} represents C_1 - C_4 -alkyl,
 in inert solvents and in the presence of a base,
 and in the case of the free hydroxyl functions ($R^4 = H$) the protective groups are removed by a customary method,
 and in the case of acids ($R^{10} = OH$), the esters are hydrolysed, and in the case of the variation of the esters ($R^{10} = OR^5$) the acids are esterified with the appropriate alcohols in the presence of a catalyst according to a customary method,
 and in the case of the amides and sulfonamides ($R^4/R^5/R^7 = -CONR^6R^9$ / $W = -CONR^{11}R^{12}$ /

-CONR¹³-SO₂R¹⁴), using amines of the formula (VIII) or sulfonamides of the formula (IX)

HN-R²⁴R²⁵ (VIII)

5 H-NR¹³-SO₂R¹⁴ (IX)

in which

R²⁴ and R²⁵ have the abovementioned meaning of R⁸, R⁹, R¹¹ and R¹²
and
10 R¹³ and R¹⁴ have the abovementioned meaning,
starting from the esters directly or starting from the free carboxylic acids, if appropriate in the
presence of above and/or an auxiliary, an amidation or sulfonamidation follows.

6. Medicaments containing at least one benzofuranyl- and -thiophenyl-alkane-carboxylic acid derivative
15 according to claim 1 to 3.

7. Medicaments according to claim 6 for the treatment and controlling acute and chronic inflammatory
processes.

20 8. Medicaments according to claim 6 for the treatment and prevention acute and chronic inflammation of
the airways, arteriosclerosis and for reducing the damage to infarct tissue after reoxygenation.

9. Use of benzofuranyl- and -thiophenyl-alkane-carboxylic acid derivatives according to claim 1 to 3 for
the production of medicaments.
25

30

35

40

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 94 10 6320

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.5)
D,X	EP-A-0 146 243 (MERCK FROSST CANADA INC.) 26 June 1985 * entire document *	1-9	C07D307/80 C07D307/81 C07D407/08 C07D405/10
A	EP-A-0 019 955 (SHELL INTERNATIONAL RESEARCH MAATSCHAPPIJ B.V.) 10 December 1980 * page 6, line 26 - page 6, line 27; claims 1-9 *	1-9	C07D405/12 C07D407/10 C07D407/12 C07D413/12 C07D413/14 A61K31/34
A	BULL. CHEM. SOC. JPN. vol. 49, no. 3, 1976 pages 737 - 740 T. HORAGUCHI ET AL. 'Furan Derivatives. II. The Cyclization of 3-(Benzofuran-3-yl)-propionic Acid Derivatives' * Scheme 3: Compound XIII; Page 739 *	1-3,5	A61K31/35 A61K31/41 A61K31/44 A61K31/445 A61K31/47 A61K31/535 C07D405/08
			TECHNICAL FIELDS SEARCHED (Int.Cl.5)
			C07D A61K
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 25 August 1994	Examiner Herz, C
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons * : member of the same patent family, corresponding document			

EPO FORM 1503 (01.82) (P04C01)